

SOLUTIONS FOR NEUTRAL AXI-DILATON GRAVITY IN 4-DIMENSIONS

T Dereli[†]

M Önder^{††}

Robin W Tucker

*School of Physics and Materials,
University of Lancaster, Bailrigg, Lancs. LA1 4YB, UK*
`r.w.tucker@lancaster.ac.uk`

ABSTRACT

We examine a 1 parameter class of actions describing the gravitational interaction between a pair of scalar fields and Einsteinian gravitation. When the parameter is positive the theory corresponds to an axi-dilatonic sector of low energy string theory. We exploit an $SL(2, \mathbb{R})$ symmetry of the theory to construct a family of electromagnetically neutral solutions with non-zero axion and dilaton charge from solutions of the Brans-Dicke theory. We also comment on solutions to the theory with negative coupling parameter.

[†] Mathematics Department, Middle East Technical University, Ankara, Turkey

^{††} Physics Department, Hacettepe University, Ankara, Turkey

1. Introduction

Among the states of low energy string theory one may identify a massless spin two graviton field in spacetime and massless spin one fields describing the Yang-Mills interactions. Besides the graviton, there arise also an antisymmetric tensor and scalar (dilaton) field with specific couplings to each other and to the graviton [1]. Depending on the topology of the compactification of internal dimensions there appear further scalar excitations with well defined coupling schemes. The pattern of all these couplings gives rise to new kinds of “duality” symmetries that may be used to discover new kinds of dilatonic black holes in 4-dimensions [2]. In the presence of the $U(1)$ gauge field such symmetries have been used to discover electrically and magnetically charged axi-dilaton “black-holes” [3] [4] [5] Such solutions tend to flat spacetime when the electric and magnetic charges tend to zero. It is of interest to enquire about the existence of electromagnetically neutral solutions with non-trivial axion and dilaton charge.

Since the above symmetries may be expressed either in terms of the geometry of the metric of the string graviton or in terms of a Weyl scaled metric (the so called Einstein frame) our discussion begins with an action that exhibits a manifest $SL(2, \mathbb{R})$ symmetry.

We examine an electrically neutral $U(1)$ sector of theory described by the spacetime classical action $S[g, \zeta] = \int \Lambda$ where

$$\Lambda = \mathcal{R} * 1 - \eta \frac{d\zeta \wedge *d\zeta^*}{q^2}. \quad (1)$$

Here $\mathcal{R}$ is the curvature scalar of the Levi-Civita connection associated with the spacetime metric g and in terms of the real scalar fields $\mathcal{A}$ and q we define the complex scalar $\zeta = \frac{\mathcal{A} + iq}{2}$. The coupling parameter η is real and denotes the ratio of the gravitational coupling constant to the scalar field interaction in the action. Expressed in terms of $\mathcal{A}$ and q

$$\Lambda = \mathcal{R} * 1 - \frac{\eta}{4q^2} dq \wedge *dq - \frac{\eta}{4q^2} d\mathcal{A} \wedge *d\mathcal{A}. \quad (2)$$

The action (1) is stationary under g and ζ variations if

$$R^{bc} \wedge *(e_a \wedge e_b \wedge e_c) + \frac{\eta}{q^2} Re(i_a d\zeta \wedge *d\zeta^* + d\zeta \wedge i_a *d\zeta^*) = 0 \quad (3)$$

$$\eta \frac{d * d\zeta}{q^2} + \frac{2i\eta}{q^3} d\zeta \wedge *d\zeta = 0 \quad (4)$$

Taking the real and imaginary parts of (4) and writing in terms of $\mathcal{A}$ and q these field equations become

$$R^{bc} \wedge *(e_a \wedge e_b \wedge e_c) + \frac{\eta}{4q^2} \tau_a[q] + \frac{\eta}{4q^2} \tau_a[\mathcal{A}] = 0 \quad (5)$$

$$d * \frac{d\mathcal{A}}{q^2} = 0 \quad (6)$$

$$d * dq - \frac{dq \wedge *dq}{q} + \frac{d\mathcal{A} \wedge *d\mathcal{A}}{q} = 0 \quad (7)$$

where the stress forms

$$\tau_a[q] = i_a dq \wedge *dq + dq \wedge i_a *dq$$

$$\tau_a[\mathcal{A}] = i_a d\mathcal{A} \wedge *d\mathcal{A} + d\mathcal{A} \wedge i_a *d\mathcal{A}.$$

We are interested in stationary spherically symmetric solutions to these equations.

2. Relation to Low Energy String States

The relation of (1) to electromagnetically neutral axion-dilaton field theory predicted by string theory can be elucidated by making a Weyl transformation on the Einstein metric: $\mathbf{g} = \lambda^2 g$ where λ is a positive scalar field. If we denote the Hodge map associated with $\mathbf{g}$ as $\star$ then for any p-form β in four dimensions one easily derives

$$*\beta = \lambda^{2p-4} \star \beta. \quad (8)$$

Thus

$$\frac{d\zeta \wedge *d\zeta^*}{q^2} = \frac{d\zeta \wedge \star d\zeta^*}{\lambda^2 q^2}. \quad (9)$$

From the definition of the curvature in terms of the metric the curvature scalar associated with $\mathbf{g}$ may be expressed as

$$\mathcal{R} * 1 = \frac{1}{\lambda^2} \mathbf{R} \star 1 + \frac{6}{\lambda^4} d\lambda \wedge \star d\lambda \quad \text{mod}(d) \quad (10)$$

If we choose $\lambda^2 = \frac{1}{q}$ then the action form is expressed as

$$\Lambda = q \mathbf{R} \star 1 - \frac{\eta - 6}{4q} dq \wedge \star dq - \frac{\eta}{4q} d\mathcal{A} \wedge \star d\mathcal{A} \quad (11)$$

which should be compared with (2). For $\eta > 0$ we may identify $\mathcal{A}$ as the axion field and ϕ as the dilaton of string theory where $q = e^{-\phi}$. In the absence of Yang-Mills gauge fields and higher order gravitational and Yang-Mills Chern-Simons couplings we may express the action in terms of the *closed* 3-form H by writing $d\mathcal{A} = e^{-\phi} \star H$. In terms of ϕ and H the effective string action form is then

$$\Lambda = e^{-\phi} \{ \mathbf{R} \star 1 - \alpha d\phi \wedge \star d\phi + \beta H \wedge \star H \} \quad (12)$$

where $\alpha = \frac{\eta-6}{4}$ and $\beta = \frac{\eta}{4}$ so $\beta - \alpha = \frac{3}{2}$.

3. Consistency with Variations of the String Action

Before seeking solutions to (5), (6) and (7) we must verify that such field equations can be derived from the low energy string action expressed in terms of the *closed* form H in order to justify the claim that the theories correspond under a Weyl scaling. In order to accommodate the closure condition on H we vary the action $S[g, q, H, \mu] = \int \Lambda'$ where

$$\Lambda' = \mathcal{R} * 1 - \frac{\eta}{4q^2} dq \wedge \star dq - \frac{\eta}{4} q^2 H \wedge \star H + \mu dH. \quad (13)$$

Since

$$\mu dH = H \wedge d\mu \quad \text{mod}(d) \quad (14)$$

the last term in (13) imposes the closure constraint [6]. This action is stationary under variations of g, q, H and μ if

$$R^{bc} \wedge *(e_a \wedge e_b \wedge e_c) + \frac{\eta}{4q^2} \tau_a[q] + \frac{\eta q^2}{4} \tau_a[H] = 0 \quad (15)$$

$$\frac{\eta}{2}d\left(\frac{*dq}{q^2}\right) + \frac{\eta}{2q^3}dq \wedge *dq - \frac{\eta}{2}H \wedge *H = 0 \quad (16)$$

$$\frac{\eta}{2}q^2 * H = d(\mu) \quad (17)$$

$$dH = 0. \quad (18)$$

Introducing the axion field $\mathcal{A}$ with the definition

$$q^2 * H = d\mathcal{A} \quad (19)$$

implies

$$d * (q^2 H) = 0 \quad (20)$$

consistent with (17) while (18) implies (6) . With the aid of the relations

$$\begin{aligned} \frac{\eta q^2}{4}\tau_a[H] &\equiv \frac{\eta q^2}{4}\{i_a H \wedge *H + H \wedge i_a *H\} \\ &= \frac{\eta}{4q^2}i_a d\mathcal{A} \wedge *d\mathcal{A} + d\mathcal{A} \wedge i_a *d\mathcal{A} = \frac{\eta}{4q^2}\tau_a[\mathcal{A}] \end{aligned} \quad (21)$$

$$H \wedge *H = -\frac{1}{q^4}d\mathcal{A} \wedge *d\mathcal{A}$$

the field equations reduce to (5) (6) and (7) as desired. Furthermore using (14) and $d(\mu) = \frac{\eta}{2}q^2 * H$ in (13) we find

$$\Lambda' = \mathcal{R} * 1 - \frac{\eta}{4q^2}dq \wedge *dq + \frac{\eta}{4}q^2 H \wedge *H \quad \text{mod}(d) \quad (22)$$

(note the sign of the third term in (13)). Substituting the relations $\frac{d\mathcal{A}}{q^2} = *H$ and $\frac{*d\mathcal{A}}{q^2} = H$ in this we recover the action form (1) .

4. Solutions for $\eta > 0$

The action defined by (1) is invariant under

$$g \mapsto g \quad (23)$$

$$\zeta \mapsto \frac{c_1\zeta + c_2}{c_3\zeta + c_4} \quad (24)$$

where the constants c_j satisfy $c_1c_4 - c_2c_3 = 1$. If one takes ζ real then the theory can be mapped by a Weyl transformation to the Brans-Dicke theory [7]. Consequently any solution of that theory can be used as a representative of the class of solutions that lie on the $SL(2, R)$ orbit of solutions solving (5) (6) and (7). For $\eta > 0$ such an orbit will then give a class of axi-dilatonic solutions with zero electromagnetic charge. Of particular interest is the orbit described by the metric

$$g = -\left(\frac{\xi + 1}{\xi - 1}\right)^{2a_1} dt \otimes dt + (\xi^2 - 1)^2 \left(\frac{\xi - 1}{\xi + 1}\right)^{2a_1} g_3 \quad (25)$$

with dilaton

$$q = \left(\frac{\xi + 1}{\xi - 1}\right)^{a_2} \quad (26)$$

and axion $\mathcal{A} = 0$. Here we employ isotropic coordinates t, r, θ, ψ and g_3 denotes the standard metric on $\mathbf{R}^3$ in terms of spherical polars r, θ, ψ . This is a particular solution if the real constants a_1, a_2 are constrained by the relation $a_1^2 + \eta a_2^2 = 1$ and $\xi \equiv -\frac{m}{2r}$ where m is a real constant. The case $a_2 = 0, a_1 = 1$ corresponds to the Schwarzschild solution with mass m . Writing $\phi = -\ln(q)$ we have for $\|\xi\| \ll 1$

$$\phi = a_2 \ln\left(\frac{\xi - 1}{\xi + 1}\right) \simeq \frac{a_2 m}{r} + \dots \quad (27)$$

and we identify the dilatonic charge of this solution to be $a_2 m$. New solutions with a non-trivial axion field can now be generated with the transformation (24). In particular with $c_1 = \cos \Theta, c_2 = \sin \Theta, c_3 = -\sin \Theta, c_4 = \cos \Theta$ one finds

$$q' = \frac{q}{\cos^2 \Theta + \sin^2 \Theta (\frac{q}{2})^2} \quad (28)$$

$$\mathcal{A}' = \frac{2 \sin \Theta \cos \Theta ((1 - \frac{q}{2})^2)}{\cos^2 \Theta + \sin^2 \Theta (\frac{q}{2})^2} \quad (29)$$

and a corresponding

$$*H' = \frac{1}{2} \sin 2\Theta d\phi. \quad (30)$$

This solution has dilaton charge $a_2 m \cos 2\Theta$ and axion charge $a_2 m \sin 2\Theta$. The purely dilatonic $U(1)$ charged solutions in references [8], [9] lie on orbits that intersect this orbit of neutral axi-dilaton solutions when the electric (and magnetic) charge is zero.

5. Solutions for $\eta < 0$

It is of interest to note that for $\eta < 0$ one may find a non-trivial $SL(2, R)$ orbit of solutions based on the metric

$$g = -e^{-\frac{2p_1}{r}} dt \otimes dt + e^{\frac{2p_1}{r}} g_3 \quad (31)$$

with

$$q = e^{-\frac{p_2}{r}} \quad (32)$$

$$\mathcal{A} = 0 \quad (33)$$

provided the real constants p_1, p_2 satisfy

$$p_1^2 + \eta p_2^2 = 0.$$

This particular solution has dilatonic charge p_2 . Under (24) with real c_j one generates the axion field

$$\mathcal{A}' = \frac{8c_2c_4 + 2c_1c_3 e^{-2\frac{p_2}{r}}}{4c_4^2 + c_3^2 e^{-2\frac{p_2}{r}}} \quad (34)$$

with axion charge

$$16p_2 \frac{2c_3c_4}{(4c_4^2 + c_3^2)^2}$$

and dilaton field

$$q' = \frac{4e^{-\frac{p_2}{r}}}{4c_4^2 + c_3^2 e^{-2\frac{p_2}{r}}} \quad (35)$$

with dilaton charge

$$4p_2 \frac{c_3^2 - c_4^2}{(4c_4^2 + c_3^2)^2}$$

The corresponding H field is

$$*H' = -c_3 c_4 d\phi. \quad (36)$$

An interesting feature of this solution is that the curvature invariant

$$*(R^{ab} \wedge *R_{ab}) = \frac{2 e^{-4p_1/r} p_1^2 (12r^2 - 16p_1 r + 2p_1^2)}{r^8}$$

is bounded in the region $r \geq 0$ with $p_1 > 0$.

6. Conclusion

We have constructed a class of electromagnetically neutral solutions to the axi-dilatonic sector of low energy string theory by exploiting a series of Weyl transformations and a manifest $SL(2, R)$ symmetry of the theory. The theory has been formulated in the “Einstein frame” with an arbitrary coupling parameter and solutions examined for both positive and negative values of this coupling parameter. We note that the singularity structure of these solutions in terms of the Levi-Civita curvature of the Einstein metric may not coincide with the corresponding curvature associated with the metric scaled by the dilaton solution.

7. Acknowledgment

RWT and TD are grateful to EPSRC for providing facilities at the 15-th UK Institute for Theoretical High Energy Physicists at the University of Southampton where this work was begun and to the Human Capital and Mobility Programme of the European Union for partial support. TD is grateful to the School of Physics and Materials, University of Lancaster, UK for its hospitality.

REFERENCES

1. E S Fradkin, A A Tseytlin, Nucl. Phys **261** (1985) 1
M Dine, N Seiberg, Phys. Rev. Letts. **55** (1985) 366
C G Callan, D Friedan, E J Martinec, M J Perry. Nucl. Phys. **B262** (1985) 593
2. A Shapere, S Trivedi, F Wilczek, Mod. Phys. Letts. **A6** (9191) 2677
3. G W Gibbons, Nucl. Phys **207** (1982) 337
G W Gibbons, K Maeda Nucl. Phys **298** (1988) 741
D Garfinkle, G T Horowitz, A Strominger, Phys. Rev. **D43** (1991) 3140
4. B A Campbell, M J Duncan, N Kaloper, K A Olive, Phys. Letts. **B251** (1990) 34
B A Campbell, N Kaloper, K A Olive, Phys. Letts. **B263** (1991) 364
R Kallosh, T Ortín Phys. Rev. **D48** (1992) 1006
S Mignemi, N R Stewert, Phys. Letts. **B298** (1993) 299
A Sen Phys. Letts. **B303** (1993) 22
5. R Kallosh, A Linde, T Ortín A Peet, A van Proeyen Phys. Rev. **D46** (1992) 5278
R Kallosh, A Peet, Phys. Rev. **D46** (1992) R5223
T Ortín Phys. Rev. **D47** (1993) 3136
R Kallosh, D Kastor, T Ortín, T Torma, preprint (1994)
6. T Dereli, R W Tucker Class. Quan. Grav. **4** (1987) 791
7. C H Brans, Phys. Rev. **125** (1962) 2194
A I Janis, E T Newman, J Winicour, Phys. Rev. Letts. **20** (1968) 878
A I Janis, D C Robinson J Winicour, Phys. Rev. **168** (1969) 1729
8. A G Agnese, M La Camera, Phys. Rev. **D49** (1994) 2126
9. M Rakhmanov, Phys. Rev. **D50** (1994) 5155